

Recoil Interaction Between Photons and The Electrons In The Plasma Of Intergalactic Space Leading To The Hubble Constant And CMB

Lyndon Ashmore

webmaster@lyndonashmore.com

ABSTRACT:

The Hubble diagram for type Ia Supernovae gives the value of the Hubble constant, H as 64 ± 3 km/s Mpc⁻¹, which, in SI units, is equal to 'hr_e/m_e per cubic metre of space' (2.1×10^{-18} s⁻¹). This coincidence could suggest a relationship between H and the electrons in the plasma of intergalactic space that act collectively and oscillate if displaced. The possibility that light from distant galaxies is absorbed and reemitted by the electrons is considered with the electron recoiling on both occasions. A double Mössbauer effect leads to a redshift in the transmitted light. Introduction of the photoabsorption cross section $2r_e\lambda$ leads to the relationship $H = 2n_e h r_e / m_e$ giving $H \approx 12$ km/s Mpc⁻¹ when n_e has the reported value of $n_e \approx 0.1 \text{ m}^{-3}$. The small amount of energy transferred to the electron by recoil is radiated as bremsstrahlung with a wavelength in the microwave region.

Key Words: Hubble constant, Intergalactic Plasma, CMB, Redshift.

Subject headings: Cosmic microwave background --- Cosmology: Galaxies: distances and redshifts --- Intergalactic medium

1. The Paradox Of The Hubble Constant and The Electron.

Whilst the conventional interpretation of cosmological redshifts is an expanding universe, some have expressed doubts that this is caused by expansion alone [1, 2, 3, 4, 5, 6]. Marmet [7] proposed a recoil interaction between photons of light and the hydrogen atoms in Intergalactic (IG) space but this would seem to have problems when one considers the discrete nature in which atoms absorb and reemit photons. However, none have reported the remarkable coincidence between the Hubble constant and the parameters of the electron ($H = h r_e / m_e$ per cubic metre of space). Nor, until now, has anyone derived a possible relationship between the two.

The Hubble diagram for type Ia Supernovae gives the value of the Hubble constant, H as 64 ± 3 km/s Mpc⁻¹ or $(2.07 \pm 0.1) \times 10^{-18}$ s⁻¹ [8]. The quantity 'hr_e/m_e' where h is the Planck constant (6.626×10^{-34} Js), r_e is the classical electron radius (2.818×10^{-15} m) and m_e is the electron rest mass (9.109×10^{-31} kg) is equal to $(2.05 \times 10^{-18} \text{ m}^3 \text{ s}^{-1})$ and so 'hr_e/m_e per cubic metre of space' has the same magnitude and dimensions as the Hubble constant. The HST key Project result for H of 72 ± 8 km/s per Mpc [9] gives a range of

$(2.1 - 2.6)\text{exp}(-18)\text{s}^{-1}$ is remarkably close to hr/m 'per cubic metre of space' when one considers that, if we are to believe in an expanding Universe, H could have had any value from zero up to the speed of light and is not supposed to be related to the electron. We must ask the question, "why is the measured value of H so close to a simple combination of the parameters of the electron if they are not related?"

These are not isolated results. Table 1 shows recent experimental values of the Hubble constant, H as selected by the ADS database. To select an unbiased sample the words "Hubble AND constant AND measurements" were fed into the database and 'return 100 items' chosen. Of these, all the papers giving an actual value for H were selected and should include the most recent results. The results for H are given in terms of hr/m per cubic metre of space. To do this the symbol k was assigned to represent the constant 'hr/m per cubic metre of space'.

| Author | Date | Bib. Code | Method Used | Value of H in terms of hr/m |
|---------------------------|---------|----------------------|----------------|-----------------------------|
| Cardone et al. | 00/2003 | 2003acfp.conf..423C | Grav. Lens | 0.91k |
| Freedman et al. | 00/2003 | 2003dhst.symp..214F | HST - Cepheids | 1.1k |
| Tikhonov et al. | 07/2002 | 2002Ap...45...253T | HST - Stars | 1.2k |
| Garinge et al. | 06/2002 | 2002MNRAS.333..318G | Xray emission | 0.89k |
| Tutui et al. | 10/2001 | 2001PASJ..53..701T | CO line T-F | 0.94k |
| Freedman et al. | 05/2001 | 2001ApJ..553..47F | HST Cepheids | 1.1k |
| Itoh et al. | 05/2001 | 2001AstHe.94.214I | Xray emission | 0.94k |
| Jensen et al. | 04/2001 | 2001ApJ.550..503J | SBF | 1.2k |
| Willick et al. | 02/2001 | 2001ApJ.548..564W | HST Cepheids | 1.3k |
| Koopmans et al | 00/2001 | 2001PASA..18..179K | Grav. lens | (0.94 - 1.1)k |
| Mauskopf et al. | 08/2000 | 2000ApJ..538..505M | Xray emission | 0.92k |
| Sakai et al. | 02/2000 | 2000ApJ..529..698S | HST Cepheids | 1.1k |
| Tanvir et al. | 11/1999 | 1999MNRAS.310..175T | HST Cepheids | 1.0k |
| Tripp et al. | 11/1999 | 1999ApJ..525..209T | Ia Supernovae | 0.97k |
| Jha et al. | 11/1999 | 1999ApJS..125..73J | Ia Supernovae | 1.0k |
| Suntzeff et al | 03/1999 | 1999AJ..117.1175S | Ia Supernova | 1.0k |
| Iwamoto et al. | 00/1999 | 1999IAUS..183..681 | Ia Supernovae | 1.0k |
| Mason et al. | 00/1999 | 1999PhDT...29M | Xray emission | 1.1k |
| Schaefer et al. | 12/1998 | 1998ApJ..509..80S | Ia Supernovae | 0.86k |
| Jha et al. | 12/1998 | 1998AAS..19310604J | Ia Supernovae | 1.0k |
| Patural et al. | 11/1998 | 1998A&A..339..671P | HIPPARCOS | 0.94k |
| Wantanabe etal | 08/1998 | 1998ApJ..503..553W | Galaxies T-F | 1.0k |
| Salaris et al. | 07/1998 | 1998MNRAS..298..166S | TRGB | 0.94k |
| Hughes et al. | 07/1998 | 1998ApJ..501..1H | Xray emission | (0.66 - 0.95)k |
| Cen et al. | 05/1998 | 1998ApJ..498L..99C | Xray emission | (0.94 - 1.3)k |
| Lauer et al | 05/1998 | 1998ApJ..449..577L | HST SBF | 1.4k |
| Average of all the values | | | | 1.0k |

Table 1.

The average of all the results was then taken. It should be noted that uncertainties were not taken into account and for those papers giving a range of values for H the middle value was taken. The average of all these results for H found by several different

techniques is equal to 1.0k i.e. hr/m per cubic metre of space. It is therefore proposed that this relationship between the Hubble constant and the electron is not a chance event.

2. THE MEDIUM WITH WHICH LIGHT INTERACTS.

This coincidence could suggest a relationship between H and the electrons in the plasma of IG space, $n_e \approx 0.1\text{m}^{-3}$ [10]. Electrons in the plasma interact simultaneously with other electrons by means of long-range Coulomb forces giving rise to a collective behaviour. Significantly, a displaced electron in the plasma of IG space will perform Simple Harmonic Motion [12] and a system of electrons that is able to oscillate is able to absorb and emit electromagnetic radiation. It is possible that photons from distant galaxies could interact with these electrons.

3. THE PROPOSED MECHANISM BY WHICH REDSHIFT OCCURS.

When photons travel through any transparent medium they are continually absorbed and reemitted by the electrons in the medium. French [13] states "the propagation of light through a medium (even a transparent one) involves a continual process of absorption of the incident light and its reemission as secondary radiation by the medium." Feynman [14] describes the transmission of light through a transparent medium simply as "photons do nothing but go from one electron to another, and reflection and transmission are really the result of an electron picking up a photon, "scratching its head", so to speak, and emitting a *new* photon." The plasma of Intergalactic space acts as a transparent medium and photons of light, as they travel through space, will be absorbed and reemitted by the electrons in this plasma. Since there is a delay at each interaction where the momentum of the photon is transferred to the electrons, the electron will recoil both on absorption and reemission - resulting in inelastic collisions [15].

A double Mössbauer effect will occur during each interaction between photon and electron. Some of the energy of the photon will be transferred to the electron and since the energy of the photon has been reduced, the frequency will reduce and the wavelength will increase. It will have undergone a 'red shift'.

Energy lost to an electron [16] during emission or absorption = $Q^2/2m_e c^2$.

Where Q is the energy of the incoming photon (hc/λ), m_e the rest mass of the electron and c , the speed of light.

This must be applied twice for absorption and reemission.

Hence, total energy lost by photon = $Q^2/m_e c^2 = h^2/\lambda^2 m_e$

(energy before interaction) - (energy after) = $h^2/\lambda^2 m_e$

$$hc/\lambda - hc/\lambda' = h^2/\lambda^2 m_e$$

λ = initial wavelength of photon, λ' = wavelength of the reemitted photon.

Multiplying through by $\lambda^2 \lambda' m_e$ and dividing by h , gives:

$$\lambda \lambda' m_e c - \lambda^2 m_e c = h \lambda'$$

Increase in wavelength, $\delta\lambda = \lambda' - \lambda$, so:

$$\lambda(\delta\lambda + \lambda)m_e c - \lambda^2 m_e c = h(\delta\lambda + \lambda)$$

$$\Rightarrow \lambda m_e c \delta\lambda + \lambda^2 m_e c - \lambda^2 m_e c = h\delta\lambda + h\lambda$$

$$\Rightarrow \delta\lambda(\lambda m_e c - h) = h\lambda$$

since $h \ll \lambda m_e c$

$$\delta\lambda = h/m_e c$$

On their journey through IG space, the photons will make many such collisions and undergo an increase in wavelength of $h/m_e c$ each time. On this basis red shift becomes a distance indicator and the distance - red shift relation becomes: photons of light from galaxies twice as far away will travel twice as far through the IG medium, make twice as many collisions and thus undergo twice the red shift.

Conservation of linear momentum will ensure the linear propagation of light.

4. THE HUBBLE LAW.

The process whereby a photon interacts with an electron and gives all its energy to the electron is known as photoabsorption and the photoabsorption cross section, σ is known from the interaction of low-energy x rays with matter [17, 18, 19].

$$\sigma = 2r_e \lambda f_2$$

Where f_2 is one of two semi-empirical atomic scattering factors depending, amongst other things, on the number of electrons in the atom. For 10 keV to 30 keV X-rays interacting with Hydrogen, f_2 has values approximately between 0 and 1. 'One' meaning

that the photon has been absorbed and the electron remaining in an excited state and 'zero' meaning that the photon was absorbed and an identical photon reemitted [13]. Since the photon frequency of light from distant galaxies is far removed from the resonant frequency of the electrons in the plasma of IG space, the photons will always be reemitted. The collision cross section for the recoil interaction considered here is therefore, $2r_e\lambda$ since f_2 only 'modulates' $2r_e\lambda$ for the atom.

On their journey through the IG medium, photons of radiation at the red end of the spectrum will encounter more collisions than photons at the blue end of the spectrum and thus undergo a greater total shift in wavelength. For a particular source, the ratio $\Delta\lambda/\lambda$ will be constant. The collision cross section for a particular photon will not be constant but will increase everytime it interacts with an electron. The photon travels shorter and shorter distances between collisions as it travels further and further and it is this that makes the red shift relation go non-linear for large red shifts.

If the initial wavelength is λ , then it will be $(\lambda + h/m_e c)$ after one collision, $(\lambda + 2h/m_e c)$ after two collisions, $(\lambda + 3h/m_e c)$ after three collisions and so on.

The mean free path of a photon in the plasma of IG space is given by $(n_e\sigma)^{-1}$ or $(2n_e r_e \lambda)^{-1}$ since $\sigma = 2r_e\lambda$.

If the photon makes a total of N collisions in travelling a distance d:

Sum of all mean free paths = d

$$\{2n_e r_e \lambda\}^{-1} + \{2n_e r_e (\lambda + h/m_e c)\}^{-1} + \{2n_e r_e (\lambda + 2h/m_e c)\}^{-1} + \{2n_e r_e (\lambda + 3h/m_e c)\}^{-1} + \dots + \{2n_e r_e (\lambda + [N-1]h/m_e c)\}^{-1} = d$$

$$\{\lambda\}^{-1} + \{\lambda + h/m_e c\}^{-1} + \{\lambda + 2h/m_e c\}^{-1} + \{\lambda + 3h/m_e c\}^{-1} + \dots + \{\lambda + [N-1]h/m_e c\}^{-1} = 2n_e r_e d$$

$$\sum_{x=0}^{N-1} \left\{ \lambda + x \left(\frac{h}{m_e c} \right) \right\}^{-1} = 2n_e r_e d$$

Since N is large and $h/m_e c$ is small ($2.43 \times 10^{-12}m$), this approximates to:

$$\int_0^{N-1} \left\{ \lambda + x \left(\frac{h}{m_e c} \right) \right\}^{-1} dx = 2n_e r_e d$$

or $1 + h\{N - 1\}/m_e c \lambda = \exp(2n_e h r_e d / m_e c)$

$$N = \lambda \exp(2n_e h r_e d / m_e c) (h / m_e c)^{-1} + 1 - \lambda (h / m_e c)^{-1} \dots \dots \dots (1)$$

The total increase in wavelength, $\Delta\lambda = N\delta\lambda$, or $Nh/m_e c$

$$\Delta\lambda = \lambda \exp(2n_e h r_e d / m_e c) + h / m_e c - \lambda$$

The red shift, z is defined as $z = \Delta\lambda / \lambda$

$$\Rightarrow z = \exp(2n_e h r_e d / m_e c) + h / m_e c \lambda - 1$$

Since $h / m_e c \lambda (= 2.42 \times 10^{-12} \lambda^{-1})$ is small for all wavelengths below X-ray,

$$z = \exp(2n_e h r_e d / m_e c) - 1$$

Using the power expansion of the exponential i.e.

$$e^x = 1 + x + x^2/2! + x^3/3! + \dots$$

Gives:

$$z = (2n_e h r_e / m_e c)d + (2n_e h r_e / m_e c)^2 d^2 / 2! + \dots (2n_e h r_e / m_e c)^3 d^3 / 3! + (2n_e h r_e / m_e c)^4 d^4 / 4! + \dots$$

In Hubble's Law, the radial velocity, v , is given as:

$$v = cz$$

$$v = c(2n_e h r_e / m_e c)d + c(2n_e h r_e / m_e c)^2 d^2 / 2! + c(2n_e h r_e / m_e c)^3 d^3 / 3! + c(2n_e h r_e / m_e c)^4 d^4 / 4! + \dots$$

Since $2n_e h r_e / m_e c$ is very small, the terms involving powers of two and above can be ignored until d becomes very large i.e. for nearby galaxies, the expression approximates to

$$v = (2n_e h r_e / m_e c)d$$

and, as $v = Hd$ (H = the Hubble constant) comparing the two equations gives

$$H = 2n_e h r_e / m_e \dots \dots \dots (2)$$

Consequently we have:

$$v = Hd + H^2 d^2 / 2! c + H^3 d^3 / 3! c^2 + H^4 d^4 / 4! c^3 + \dots$$

i.e. $v = c \{ \exp(Hd/c) - 1 \}$ and

$$z = \exp(Hd/c) - 1$$

It should be noted that this relationship between redshift, z and distance, d is identical to that first proposed by Zwicky in 1929 [20].

5. COMPARISON WITH EXPERIMENTAL RESULTS.

This theory predicts (equation 2) $H = 2n_e h r_e / m_e$

or $H = 4.10 \times 10^{-18} n_e \text{ s}^{-1}$

As cited earlier, $n_e \approx 0.1 \text{ m}^{-3}$ [10,11] and using this value to predict H gives:

$$H \approx 0.41 \times 10^{-18} \text{ s}^{-1} \quad (H \approx 12 \text{ km/s Mpc}^{-1})$$

This is in good agreement with the experimental values ([8] i.e. $H = (2.1 \pm 0.1) \times 10^{-18} \text{ s}^{-1}$). To predict the experimentally derived value of H of $2.1 \times 10^{-18} \text{ s}^{-1}$ ($64 \pm 3 \text{ km/s Mpc}$) requires $n_e \approx 0.5 \text{ m}^{-3}$ compared to the cited value of $n_e \approx 0.1 \text{ m}^{-3}$. Light of wavelength $5 \times 10^{-7} \text{ m}$ would have a collision cross-section of $2.8 \times 10^{-21} \text{ m}^2$ and each photon would, on average, make one collision with an electron in the plasma of IG space every 75,000 light year.

Published statistical tests on redshift data show that the Hubble diagram is straight up to $z \approx 0.1$, goes nonlinear at $z \approx 0.8$, is quadratic at $z \approx 2.8 - 3.6$ and for redshifts above this, follows a non simple power law ([8, 21, 22, 23]. However, it has recently been shown [24] that data from the Calan/Tololo Supernova survey can verify this exponential law with a value of H of 72 km/s per Mpc ie $1.13 h r_e / m_e \text{ per m}^3$ ($n_e = 0.57 \text{ m}^{-3}$) if the data is not 'corrected' for the relativistic effects of expansion first.




Fig 1

That is the data fits this theory's predicted exponential Hubble law provided that we do not assume that the Universe is expanding and manipulate the data accordingly. This theory's predicted exponential Hubble curve is shown in Fig 1 for comparison.

6. COSMIC MICROWAVE BACKGROUND (CMB).

The recoiling electron will be brought to rest by Coulomb interactions with all the electrons contained within a Debye sphere of radius λ_D . The decelerating electron will emit transmission radiation (TR) i.e. bremsstrahlung. There are two emission channels of the system, 'intrinsic emission' by the decelerating electron, and 'emission by the medium' where the background electrons radiate energy.

Intrinsic radiation arises when the recoiling electron exchanges a virtual photon with the external field (set up by the large number of coulomb centers) with momentum \mathbf{q} and emits a quantum with momentum \mathbf{k} . The medium or external field in which the recoiling electron is moving radiates when the virtual photon of momentum \mathbf{q} results in the production of radiation by background electrons contained within the Debye sphere [25].

The interactions between light and the electrons are non-relativistic and the initial and final states of the electron belong to the continuous spectrum. The photon frequency of the transmission radiation f_{cmb} is given by:

$$hf_{\text{cmb}} = (1/2m_e)(p^2 - p'^2)$$

Where $p = m_e v$ and $p' = m_e v'$ are the initial and final momentum of the electron [26].

The electron returns to rest after absorption and reemission and so the wavelength of the transmission radiation λ_{CMB} is given by:

$$\lambda_{\text{CMB}} = 2m_e \lambda^2 c / h$$

Light of wavelength $5 \times 10^{-7} \text{m}$ gives rise to TR of wavelength 0.21m . In IG space, the dominant background photons are microwaves, having peak energy of $6 \times 10^{-4} \text{eV}$ and a photon density of about 400 per cm^{-3} [27, 28]. In this theory, these background photons ($\lambda = 2.1 \times 10^{-3} \text{m}$) would be given off as TR by a photon of wavelength $5 \times 10^{-8} \text{m}$ (i.e. Ultra Violet radiation) interacting with an electron.

7. DISCUSSION.

This proposed theory has successes in predicting values of H and λ_{CMB} that have the same magnitude as experimental values. As to whether this proposed interaction makes up the whole of the redshift or just a part of it will not be known until n_e has been determined to a greater accuracy. It also shows a relationship between H and 'hr_e/m_e per cubic metre' which could explain the remarkable coincidence between their magnitudes. As scientists, we must always be suspicious of quantities that are equal but do not appear to be related. The theory still has to explain the 'surface brightness test' [29] and the time dilation in supernova light curves [30, 31, 32, 33, 34].

However, the value of H quoted here (64 ± 3 km/s Mpc⁻¹) is only one value and other techniques and other workers give differing values. A value of 70 ± 7 km/s Mpc⁻¹ can be said to represent present data from all areas [35] and thus all agree that values of H lie in the range 1.0 to 1.2 times 'hr_e/m_e per cubic metre of space'. With the 'big bang' theory, H could have had any initial value (as far as we know) and the effects of gravity and 'vacuum energy' make H time dependent, changing in magnitude from this original value. How probable is it that the first time we measure H with some accuracy it has the same value as 'hr_e/m_e per unit volume of space' (especially when these constants carry such importance in the scattering of light). To complicate matters further, the age of the Universe is often quoted as H^{-1} which we now realize to be equal to between 0.8 and 1.0 times m_e/hr_e - this is a test that the expanding universe 'fails' and now has to explain.

Submitted and registered for publication, March 2003.

Accepted for Publication June 2004.

REFERENCES:

1. Burbidge. Et al. 1971 ApJ, 170, 233 - 240
2. Marič, Z., Moles, M., and Vigier, J. P. 1977, Lett. Nuovo Cimento, 18, 269 - 276
3. Arp, H. 1987 Quasars, Redshifts, and Controversies. (Interstellar Media, Berkeley)
4. Cohen et al. 1988 Ap.J 329 1 - 7
5. Arp, H. 1990. Nature, 346, 807 - 812
6. Arp, H. ApJ, 1994 430, 74 - 82,
7. Marmet, P. 1988 Physics Essays, Vol 1No.1 p24-32
8. Reiss, Press, Kirshner. 1996 ApJ. 473. 88 - 109
9. Freedman, W. et al 2003 dhst.symp..214 - 222F
10. Peebles, P.J.E. 1993 Principles of Physical Cosmology
11. Deffayet, C., Harari, D., Uzan, JP., Zaldarriaga, M. 2002

- PhRv. D 66d3517D 6 pages.
12. Mitchner, M., Kruger, C.H. Partially Ionized gases 1973 (Wiley, p138.USA,)
 13. French.A.P. 1968a Special relativity (p128. Nelson. London)
 14. Feynman.R."Q.E.D.- the strange story of light and matter".P76. Penguin.London.1990.
 15. Berestetskii, V.B., Lifshitz,E.M. Pitaevskii,L.P. 1982a Quantum Electrodynamics Volume 4, second edition. (p161 & 221. Butterworth Heinemann, Oxford. UK.)
 16. French.A.P. 1968b Special relativity (p176-182. Nelson. London)
 17. Henke, B.L., Gullikson, E.M., Davis, J.C. 1993 Atomic Data and Nuclear Data Tables, 54, p181-342
 18. Henke, B.L., Gullikson, E.M., Davis, J.C. 2001 X-Ray Data Booklet chap 1 p 44/52(LBNL/PUB-490 Rev. 2 Lawrence Berkeley National Laboratory, University of California, Berkely, CA 94720)(also at: http://www-cxro.lbl.gov/optical_constants/intro.html)
 19. Hubbell, J.H.,Veigele, W.J., Briggs, E.A., Brown, R.T., Cromer, D.T. Howerton, R.J. 1975 J. Phys. Chem. Ref. Data 4, 471-538; 1977 erratum in 6, 615-616
 20. Zwicky, F. 1929 Proc. Nat. Acad. Sci., 773 - 785
 21. Sandage,Kristian,Westphal. 1976 ApJ. 205. 688-695
 22. Segal.I.E. MNRAS 1989. 237, 17 - 37
 23. Soneira.R.M. 1979 ApJ. 230. L63 -L65
 24. Khaidarov, K. <http://bourabai.narod.ru/universum.htm>
 25. Platonov, K. Yu. Fleishman, G.D. 2002 Uspekhi Fizicheskikh Nauk 172 (3) 241 - 300
 26. Berestetskii, V.B., Lifshitz,E.M. Pitaevskii,L.P. 1982b Quantum Electrodynamics (Volume 4, second edition. p389. Butterworth Heinemann, Oxford. UK.).
 27. Peebles, P.J.E., Schramm, D.N. Kron, R.G., Yurner, E.L. 1991 Nature vol 352. 769 - 776
 28. Nagano, M. & Watson, A.A. July 2000 Reviews of Modern Physics, Vol 72 No. 3, 689 - 732 American Physical soc
 29. Sandage, A. 1991 ApJ 370 455 - 473
 30. Reiss,A.G. Filippenko, A.V. Leonard, D.C. 1997, Astronomical journal, 114,722-729
 31. Perlmutter et. al. Nature 1998. 391 51 - 54
 32. Perlmutter, S. et al., 1999 ApJ 517, 565 - 586
 33. Riess, A.G. et al., 1998 AJ 116, 1009 - 1038.
 34. Reiss, A.G. et al., 2001 ibid. 560, 49.
 35. Krishner,R.P. 2000 The Extravagant Universe" (p98. Princeton University Press. Princeton. N.J.)